

Table 2. Mean abundance by station grouping of those species that contribute most to the within group similarity from the first ordination (see text). Highest mean abundances are highlighted.

Species	Station group					
	1	2	3	4	5	6
<i>Euphausia superba</i>	0.51	1.66	1.53	1.97	2.56	1.13
<i>Metridia</i> spp.	9.68	6.92	4.98	7.48	0.92	5.88
<i>Ctenocalanus</i> spp.	9.62	6.74	3.15	7.10	0.61	2.83
<i>Oithona</i> spp.	9.44	7.75	2.91	2.79	0.47	1.52
<i>Calanoides acutus</i>	9.13	5.95	2.98	8.68	1.01	7.15
<i>Rhincalaus gigas</i>	7.65	4.02	2.01	3.15	0.53	4.14
<i>Calanus propinquus</i>	7.16	5.87	3.80	7.37	1.29	7.88
<i>Scolecithricella minor</i>	5.81	4.99	3.52	1.40	0.00	3.64
<i>Euchaeta antarctica</i>	5.19	5.24	3.27	3.30	0.00	4.02
<i>Thysanoessa vicina</i>	7.48	4.44	0.98	5.76	1.13	7.30
<i>Thysanoessa</i> spp. calyptopes	6.32	4.81	1.65	3.99	0.00	6.52
<i>Thysanoessa macrura</i>	2.65	3.62	3.59	2.06	4.01	1.52
<i>Euphausia superba</i> larvae	2.84	2.09	1.23	8.89	0.39	4.11
Chaetognatha	7.19	6.09	3.12	4.87	1.10	4.54
<i>Salpa thompsoni</i>	1.18	2.87	3.05	2.77	2.94	0.00
<i>Limacina helicina</i>	6.65	0.79	0.11	1.16	0.00	3.99
<i>Limacina</i> sp.	0.00	1.19	0.35	1.09	2.33	0.00